


06: Truth & Proof

Logical Foundations of Cyber-Physical Systems



André Platzer

Karlsruhe Institute of Technology
Department of Informatics

Computer Science Department
Carnegie Mellon University

1 Learning Objectives

2 Sequent Calculus

- Propositional Proof Rules
- Soundness of Proof Rules
- Proofs with Dynamics
- Contextual Equivalence
- Quantifier Proof Rules
- A Sequent Proof for Single-hop Bouncing Balls

3 Real Arithmetic

- Real Quantifier Elimination
- Instantiating Real-Arithmetic Quantifiers
- Weakening by Removing Assumptions
- Abbreviating Terms to Reduce Complexity
- Substituting Equations into Formulas
- Creatively Cutting to Transform Questions

4 Summary

1 Learning Objectives

2 Sequent Calculus


- Propositional Proof Rules
- Soundness of Proof Rules
- Proofs with Dynamics
- Contextual Equivalence
- Quantifier Proof Rules
- A Sequent Proof for Single-hop Bouncing Balls

3 Real Arithmetic

- Real Quantifier Elimination
- Instantiating Real-Arithmetic Quantifiers
- Weakening by Removing Assumptions
- Abbreviating Terms to Reduce Complexity
- Substituting Equations into Formulas
- Creatively Cutting to Transform Questions


4 Summary

systematic reasoning for CPS
verifying CPS models at scale
pragmatics: how to use axiomatics to justify truth
structure of proofs and their arithmetic



discrete+continuous relation
with evolution domains

analytic skills for CPS



Syntax defines the notation

What problems are we allowed to write down?

Semantics what carries meaning.

What real or mathematical objects does the syntax stand for?

Axiomatics internalizes semantic relations into universal syntactic transformations.

Pragmatics how to use axiomatics to justify syntactic rendition of semantical concepts. How to do a proof?

1 Learning Objectives

2 Sequent Calculus

- Propositional Proof Rules
- Soundness of Proof Rules
- Proofs with Dynamics
- Contextual Equivalence
- Quantifier Proof Rules
- A Sequent Proof for Single-hop Bouncing Balls

3 Real Arithmetic

- Real Quantifier Elimination
- Instantiating Real-Arithmetic Quantifiers
- Weakening by Removing Assumptions
- Abbreviating Terms to Reduce Complexity
- Substituting Equations into Formulas
- Creatively Cutting to Transform Questions

4 Summary

Definition (Sequent)

$$\Gamma \vdash \Delta$$

has the same meaning as $\bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q$.

The *antecedent* Γ and *succedent* Δ are finite sets of dL formulas.

Definition (Soundness of sequent calculus proof rules)

$$\frac{\Gamma_1 \vdash \Delta_1 \quad \dots \quad \Gamma_n \vdash \Delta_n}{\Gamma \vdash \Delta}$$

is *sound* iff validity of all premises implies validity of conclusion:

If $\models (\Gamma_1 \vdash \Delta_1)$ and \dots and $\models (\Gamma_n \vdash \Delta_n)$ then $\models (\Gamma \vdash \Delta)$

Definition (Sequent)

$$\Gamma \vdash \Delta$$

has the same meaning as $\bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q$.

The *antecedent* Γ and *succedent* Δ are finite sets of dL formulas.

Definition (Soundness of sequent calculus proof rules)

construct proofs up



$$\frac{\Gamma_1 \vdash \Delta_1 \quad \dots \quad \Gamma_n \vdash \Delta_n}{\Gamma \vdash \Delta}$$

is *sound* iff validity of all premises implies validity of conclusion:

If $\models (\Gamma_1 \vdash \Delta_1)$ and ... and $\models (\Gamma_n \vdash \Delta_n)$ then $\models (\Gamma \vdash \Delta)$

Definition (Sequent)

$$\Gamma \vdash \Delta$$

has the same meaning as $\bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q$.

The *antecedent* Γ and *succedent* Δ are finite sets of dL formulas.

Definition (Soundness of sequent calculus proof rules)

$$\begin{array}{ccc}
 \text{construct proofs up} \uparrow & \frac{\Gamma_1 \vdash \Delta_1 \quad \dots \quad \Gamma_n \vdash \Delta_n}{\Gamma \vdash \Delta} & \downarrow \text{validity transfers down}
 \end{array}$$

is *sound* iff validity of all premises implies validity of conclusion:

$$\text{If } \models (\Gamma_1 \vdash \Delta_1) \text{ and } \dots \text{ and } \models (\Gamma_n \vdash \Delta_n) \text{ then } \models (\Gamma \vdash \Delta)$$

$$\wedge L \frac{}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$\wedge L$: assume conjuncts separately

It successively handles all top-level \wedge in assumptions but not nested in $A \vee (B \wedge C) \vdash C$ which needs rules for other propositional operators

$$\wedge R \frac{}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$\wedge R$: prove conjuncts separately

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta} \quad \vee R \frac{}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta} \quad \vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$\vee R$: split disjunctions in succedent where comma has a disjunctive meaning

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{}{\Gamma, P \vee Q \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$\vee L$: handle disjunctive assumption by one proof for each assumed disjunct

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{}{\Gamma \vdash P \rightarrow Q, \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$\rightarrow R$: prove implication by assuming LHS when proving RHS

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{}{\Gamma, P \rightarrow Q \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$\rightarrow L$: assume RHS of an assumed implication after proving its LHS

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$\neg R$: prove $\neg P$ by proving contradiction (or Δ options) from assumption P

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$\neg L$: assume $\neg P$ by proving its opposite P

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

id: proof done (marked *) when succedent to prove is in antecedent

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

id: only way to finish a proof (in propositional logic!)

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{}{\Gamma \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta}$$

Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta}$$

cut: Show lemma C and then assume lemma C

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\top R \frac{}{\Gamma \vdash \text{true}, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta}$$

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\top R \frac{}{\Gamma \vdash \text{true}, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta}$$

$\top R$: proof done (marked *) when proving trivial *true* (used rarely)

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\top R \frac{}{\Gamma \vdash \text{true}, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta}$$

$\top R$: what rule to use when *true* in antecedent?

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\top R \frac{}{\Gamma \vdash \text{true}, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta}$$

$$\perp L \frac{}{\Gamma, \text{false} \vdash \Delta}$$

Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\top R \frac{}{\Gamma \vdash \text{true}, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta}$$

$$\perp L \frac{}{\Gamma, \text{false} \vdash \Delta}$$

$\perp L$: proof done (marked *) when assuming trivial *false* (used rarely)

\mathcal{A} Propositional Proof Rules of Sequent Calculus

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$$

$$\vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta}$$

$$\neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta}$$

$$\wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta}$$

$$\vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta}$$

$$\neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta}$$

$$\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta}$$

$$\text{id} \frac{}{\Gamma, P \vdash P, \Delta}$$

$$\top R \frac{}{\Gamma \vdash \text{true}, \Delta}$$

$$\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta}$$

$$\text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta}$$

$$\perp L \frac{}{\Gamma, \text{false} \vdash \Delta}$$

$\perp L$: what rule to use when *false* in succedent?

$$\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)$$

$$\rightarrow R \frac{\frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)}}{\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)}}$$

$$\begin{array}{c}
 \frac{\frac{\overline{v^2 \leq 10 \wedge b > 0 \vdash b > 0} \quad \overline{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0) \vee v^2 \leq 10}}{\wedge R} \quad \overline{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)}}{\rightarrow R} \\
 \vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)
 \end{array}$$

$$\begin{array}{c}
 \frac{}{\overline{v^2 \leq 10, b > 0 \vdash b > 0}} \\
 \wedge L \frac{}{\overline{v^2 \leq 10 \wedge b > 0 \vdash b > 0}} \quad \frac{}{\overline{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0) \vee v^2 \leq 10}} \\
 \wedge R \frac{}{\overline{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)}} \\
 \rightarrow R \frac{}{\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)}
 \end{array}$$



Sequent Proof Example (Simple)

$$\begin{array}{c} * \\ \text{id} \frac{}{v^2 \leq 10, b > 0 \vdash b > 0} \\ \wedge L \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0} \quad \frac{}{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0) \vee v^2 \leq 10} \\ \wedge R \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\ \rightarrow R \frac{}{\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \end{array}$$



Sequent Proof Example (Simple)

$$\begin{array}{c}
 * \\
 \text{id} \frac{}{v^2 \leq 10, b > 0 \vdash b > 0} \\
 \wedge L \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0} \quad \vee R \frac{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0), v^2 \leq 10}{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0) \vee v^2 \leq 10} \\
 \wedge R \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 \rightarrow R \frac{}{\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)}
 \end{array}$$



Sequent Proof Example (Simple)

$$\begin{array}{c}
 \text{id} \frac{}{v^2 \leq 10, b > 0 \vdash b > 0} \\
 \wedge L \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0} \\
 \wedge R \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 \rightarrow R \frac{}{\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 * \frac{}{v^2 \leq 10, b > 0 \vdash \neg(v \geq 0), v^2 \leq 10} \\
 \wedge L \frac{}{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0), v^2 \leq 10} \\
 \vee R \frac{}{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0) \vee v^2 \leq 10}
 \end{array}$$



Sequent Proof Example (Simple)

$$\begin{array}{c} \text{id} \frac{}{v^2 \leq 10, b > 0 \vdash b > 0} \\ \wedge L \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0} \\ \wedge R \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\ \rightarrow R \frac{}{\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \end{array} \quad \begin{array}{c} * \\ \text{id} \frac{}{v^2 \leq 10, b > 0 \vdash \neg(v \geq 0), v^2 \leq 10} \\ \wedge L \frac{}{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0), v^2 \leq 10} \\ \vee R \frac{}{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0) \vee v^2 \leq 10} \\ * \end{array}$$

Lemma

$$\wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta} \text{ is sound}$$

Lemma

$\wedge R$ $\frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$ is sound: conclusion valid if all premises valid.

Lemma

$\wedge R$ $\frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta}$ is sound: conclusion valid if all premises valid.

Proof

using $\llbracket P \wedge Q \rrbracket = \llbracket P \rrbracket \cap \llbracket Q \rrbracket$.

WLOG: $\omega \in \llbracket G \rrbracket$ for all $G \in \Gamma$ and $\omega \notin \llbracket D \rrbracket$ for all $D \in \Delta$ (why?)

By premise: $\omega \in \llbracket \Gamma \vdash P, \Delta \rrbracket$ and $\omega \in \llbracket \Gamma \vdash Q, \Delta \rrbracket$

By WLOG: $\omega \in \llbracket P \rrbracket$ and $\omega \in \llbracket Q \rrbracket$ □

By semantics: $\omega \in \llbracket P \wedge Q \rrbracket$

By definition: $\omega \in \llbracket \Gamma \vdash P \wedge Q, \Delta \rrbracket$

Theorem

dL sequent calculus is sound: every dL formula with a proof is valid.



Theorem

dL *sequent calculus* is sound: every dL *sequent* with a proof is valid.

Proof (by induction on structure of sequent calculus proof).



Theorem

dL *sequent calculus is sound: every dL sequent with a proof is valid.*

Proof (by induction on structure of sequent calculus proof).

- 0 Proofs without rule uses only prove dL axioms, which are sound.

Theorem

dL *sequent calculus* is sound: every dL sequent with a proof is valid.

Proof (by induction on structure of sequent calculus proof).

- 0 Proofs without rule uses only prove dL axioms, which are sound.
- 1 Sequent proof ends with some proof step:

$$\frac{\Gamma_1 \vdash \Delta_1 \quad \dots \quad \Gamma_n \vdash \Delta_n}{\Gamma \vdash \Delta}$$

The subproof of each premise $\Gamma_i \vdash \Delta_i$ is smaller, so $\models \Gamma_i \vdash \Delta_i$ by IH.
All dL proof rules are proved sound, also the one used above, i.e.:

If $\models (\Gamma_1 \vdash \Delta_1)$ and ... and $\models (\Gamma_n \vdash \Delta_n)$ then $\models (\Gamma \vdash \Delta)$

Thus, $\models (\Gamma \vdash \Delta)$. □

Theorem

dL *sequent calculus* is sound: every dL sequent with a proof is valid.

Proof (by induction on structure of sequent calculus proof).

- 0 Proofs without rule uses only prove dL axioms, which are sound.
- 1 Sequent proof ends with some proof step:

$$\frac{\Gamma_1 \vdash \Delta_1 \quad \dots \quad \Gamma_n \vdash \Delta_n}{\Gamma \vdash \Delta}$$

The subproof of each premise $\Gamma_i \vdash \Delta_i$ is smaller, so $\models \Gamma_i \vdash \Delta_i$ by IH.
All dL proof rules are proved sound, also the one used above, i.e.:

If $\models (\Gamma_1 \vdash \Delta_1)$ and ... and $\models (\Gamma_n \vdash \Delta_n)$ then $\models (\Gamma \vdash \Delta)$

Thus, $\models (\Gamma \vdash \Delta)$. □

▶ **Todo** Always make sure every axiom and proof rule we adopt is sound!

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

$$[\cup]R \frac{}{\Gamma \vdash [\alpha \cup \beta]P, \Delta}$$

$$[\cup]L \frac{}{\Gamma, [\alpha \cup \beta]P \vdash \Delta}$$

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

$$[\cup]R \frac{\Gamma \vdash [\alpha]P \wedge [\beta]P, \Delta}{\Gamma \vdash [\alpha \cup \beta]P, \Delta}$$

$$[\cup]L \frac{}{\Gamma, [\alpha \cup \beta]P \vdash \Delta}$$

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

$$[\cup]R \frac{\Gamma \vdash [\alpha]P \wedge [\beta]P, \Delta}{\Gamma \vdash [\alpha \cup \beta]P, \Delta}$$

$$[\cup]L \frac{\Gamma, [\alpha]P \wedge [\beta]P \vdash \Delta}{\Gamma, [\alpha \cup \beta]P \vdash \Delta}$$

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

$$[\cup]R \frac{\Gamma \vdash [\alpha]P \wedge [\beta]P, \Delta}{\Gamma \vdash [\alpha \cup \beta]P, \Delta} \quad \text{Boring! Already follow from the axiom}$$

$$[\cup] [\alpha \cup \beta]P \leftrightarrow [\alpha]P \wedge [\beta]P$$

$$[\cup]L \frac{\Gamma, [\alpha]P \wedge [\beta]P \vdash \Delta}{\Gamma, [\alpha \cup \beta]P \vdash \Delta}$$

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

$$[\cup]R \frac{\Gamma \vdash [\alpha]P \wedge [\beta]P, \Delta}{\Gamma \vdash [\alpha \cup \beta]P, \Delta}$$

Boring! Already follow from the axiom

$$[\cup] [\alpha \cup \beta]P \leftrightarrow [\alpha]P \wedge [\beta]P$$

$$[\cup]L \frac{\Gamma, [\alpha]P \wedge [\beta]P \vdash \Delta}{\Gamma, [\alpha \cup \beta]P \vdash \Delta}$$

Rules $[\cup]R, [\cup]L$ would only apply top-level,
not in any other logical context such as
 $[x'' = -g]_-$

$$[\cup] \frac{}{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}$$

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

$$[U]R \frac{\Gamma \vdash [\alpha]P \wedge [\beta]P, \Delta}{\Gamma \vdash [\alpha \cup \beta]P, \Delta} \quad \text{Boring! Already follow from the axiom } [U] [\alpha \cup \beta]P \leftrightarrow [\alpha]P \wedge [\beta]P$$

$$[U]L \frac{\Gamma, [\alpha]P \wedge [\beta]P \vdash \Delta}{\Gamma, [\alpha \cup \beta]P \vdash \Delta} \quad \text{Rules } [U]R, [U]L \text{ would only apply top-level, not in any other logical context such as } [x'' = -g]_-$$

Contextual Equivalence: substituting equals for equals

$$\text{CER} \frac{\Gamma \vdash C(Q), \Delta \quad \vdash P \leftrightarrow Q}{\Gamma \vdash C(P), \Delta} \quad \text{CEL} \frac{\Gamma, C(Q) \vdash \Delta \quad \vdash P \leftrightarrow Q}{\Gamma, C(P) \vdash \Delta}$$

$$[U] \frac{}{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}$$

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

$$[\cup]R \frac{\Gamma \vdash [\alpha]P \wedge [\beta]P, \Delta}{\Gamma \vdash [\alpha \cup \beta]P, \Delta}$$

Boring! Already follow from the axiom

$$[\cup] [\alpha \cup \beta]P \leftrightarrow [\alpha]P \wedge [\beta]P$$

$$[\cup]L \frac{\Gamma, [\alpha]P \wedge [\beta]P \vdash \Delta}{\Gamma, [\alpha \cup \beta]P \vdash \Delta}$$

Rules $[\cup]R, [\cup]L$ would only apply top-level,
not in any other logical context such as
 $[x'' = -g]_-$

Contextual Equivalence: substituting equals for equals

$$\text{CER} \frac{\Gamma \vdash C(Q), \Delta \quad \vdash P \leftrightarrow Q}{\Gamma \vdash C(P), \Delta}$$

$$\text{CEL} \frac{\Gamma, C(Q) \vdash \Delta \quad \vdash P \leftrightarrow Q}{\Gamma, C(P) \vdash \Delta}$$

$$[?x=0; v := -cv \cup ?x \geq 0]B(x, v) \leftrightarrow [?x=0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v)$$

$$[\cup] \frac{}{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}$$

Have: Left and right proof rule for all propositional connectives

Need: Left and right proof rule for all top-level operators in all modalities?

$$[\cup]R \frac{\Gamma \vdash [\alpha]P \wedge [\beta]P, \Delta}{\Gamma \vdash [\alpha \cup \beta]P, \Delta}$$

Boring! Already follow from the axiom

$$[\cup] [\alpha \cup \beta]P \leftrightarrow [\alpha]P \wedge [\beta]P$$

$$[\cup]L \frac{\Gamma, [\alpha]P \wedge [\beta]P \vdash \Delta}{\Gamma, [\alpha \cup \beta]P \vdash \Delta}$$

Rules $[\cup]R, [\cup]L$ would only apply top-level, not in any other logical context such as $[x'' = -g]_-$

Contextual Equivalence: substituting equals for equals

$$\text{CER} \frac{\Gamma \vdash C(Q), \Delta \quad \vdash P \leftrightarrow Q}{\Gamma \vdash C(P), \Delta}$$

$$\text{CEL} \frac{\Gamma, C(Q) \vdash \Delta \quad \vdash P \leftrightarrow Q}{\Gamma, C(P) \vdash \Delta}$$

$$[?x=0; v := -cv \cup ?x \geq 0]B(x, v) \leftrightarrow [?x=0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v)$$

$$[\cup] \frac{A \vdash [x'' = -g]([?x = 0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}$$



$$[\cdot] \frac{}{\vdash [a := -b; c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$$[a := -b; c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c)) \leftrightarrow$$

$$[a := -b][c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c)) \text{ by } [;]$$

$$\frac{[;]}{[a := -b][c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$$\frac{[;]}{[a := -b; c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$[a := -b][c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c)) \leftrightarrow$
 $[c := 10](v^2 \leq 10 \wedge -(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))$ by $[:=]$

$$\frac{}{[:=] \vdash [c := 10](v^2 \leq 10 \wedge -(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$$\frac{}{[:=] \vdash [a := -b][c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$$\frac{}{[:] \vdash [a := -b; c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$[c := 10](v^2 \leq 10 \wedge -(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c)) \leftrightarrow$
 $v^2 \leq 10 \wedge -(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)$ by $[:=]$

$$\frac{}{\vdash v^2 \leq 10 \wedge -(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)}$$

$$[:=] \frac{}{\vdash [c := 10](v^2 \leq 10 \wedge -(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$$[:=] \frac{}{\vdash [a := -b][c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$$[:] \frac{}{\vdash [a := -b; c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}$$

$$\begin{array}{c}
 \text{id} \frac{}{v^2 \leq 10, b > 0 \vdash b > 0} \quad \text{id} \frac{}{v^2 \leq 10, b > 0 \vdash \neg(v \geq 0), v^2 \leq 10} \\
 \text{\textcircled{*}} \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0} \quad \text{\textcircled{*}} \frac{}{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0), v^2 \leq 10} \\
 \text{\textcircled{\wedge}L} \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0} \quad \text{\textcircled{\wedge}R} \frac{}{v^2 \leq 10 \wedge b > 0 \vdash \neg(v \geq 0) \vee v^2 \leq 10} \\
 \text{\textcircled{\wedge}R} \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 \text{\textcircled{\rightarrow}R} \frac{}{\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 \frac{}{\vdash v^2 \leq 10 \wedge \neg(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 \text{\textcircled{[:=]}} \frac{}{\vdash [c := 10](v^2 \leq 10 \wedge \neg(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))} \\
 \text{\textcircled{[:=]}} \frac{}{\vdash [a := -b][c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))} \\
 \text{\textcircled{[;]}} \frac{}{\vdash [a := -b; c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}
 \end{array}$$

$$\begin{array}{c}
 \text{id} \frac{}{v^2 \leq 10, b > 0 \vdash b > 0} \\
 \wedge^L \frac{\text{id} \frac{}{v^2 \leq 10, b > 0 \vdash b > 0}}{v^2 \leq 10 \wedge b > 0 \vdash b > 0} \\
 \wedge^R \frac{}{v^2 \leq 10 \wedge b > 0 \vdash b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 \rightarrow R \frac{}{\vdash v^2 \leq 10 \wedge b > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 \frac{}{\vdash v^2 \leq 10 \wedge \neg(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq 10)} \\
 [:=] \frac{}{\vdash [c := 10](v^2 \leq 10 \wedge \neg(-b) > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))} \\
 [:=] \frac{}{\vdash [a := -b][c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))} \\
 [i] \frac{}{\vdash [a := -b; c := 10](v^2 \leq 10 \wedge -a > 0 \rightarrow b > 0 \wedge (\neg(v \geq 0) \vee v^2 \leq c))}
 \end{array}$$

Need to reason about real arithmetic

Here: to glue previous propositional proof with this dynamic proof

$$\forall R \frac{}{\Gamma \vdash \forall x p(x), \Delta}$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta}$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$\forall R$: show for fresh variable y about which we can't know anything

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\exists R \frac{}{\Gamma \vdash \exists x p(x), \Delta}$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta}$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e)$$

$\exists R$: enough to show for any witness term e

Quantifier Proof Rules

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\forall L \frac{}{\Gamma, \forall x p(x) \vdash \Delta}$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e)$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta}$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e)$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} \quad (\text{arbitrary term } e)$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e)$$

$\forall L$: even holds for arbitrary term e

Quantifier Proof Rules

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} \quad (\text{arbitrary term } e)$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e)$$

$$\exists L \frac{}{\Gamma, \exists x p(x) \vdash \Delta}$$

Quantifier Proof Rules

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} \quad (\text{arbitrary term } e)$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e)$$

$$\exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta}$$

Quantifier Proof Rules

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} \quad (\text{arbitrary term } e)$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e)$$

$$\exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} \quad (y \notin \Gamma, \Delta, \exists x p(x))$$

$\exists L$: assume for fresh variable y about which we can't know anything

Quantifier Proof Rules

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x))$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} \quad (\text{arbitrary term } e)$$

$$\exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e)$$

$$\exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} \quad (y \notin \Gamma, \Delta, \exists x p(x))$$

Important: soundness means that conclusion valid if all premises valid.

$$\begin{array}{c} \rightarrow^R \\ \hline \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v) \\ \quad A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0 \\ \quad B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H \\ \quad \{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\} \end{array}$$

$$\frac{[i] \overline{A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}}{\rightarrow R \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$$\begin{array}{c}
A \vdash \forall t \geq 0 \left(\left(H - \frac{g}{2}t^2 = 0 \rightarrow B\left(H - \frac{g}{2}t^2, -c(-gt)\right) \right) \wedge \left(H - \frac{g}{2}t^2 \geq 0 \rightarrow B\left(H - \frac{g}{2}t^2, -g\right) \right) \right) \\
\hline
[:=] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2] \left((x = 0 \rightarrow B(x, -c(-gt))) \wedge (x \geq 0 \rightarrow B(x, -gt)) \right) \\
\hline
[:=] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2] [v := -gt] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
\hline
[.] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2; v := -gt] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
\hline
['] A \vdash [x'' = -g] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
\hline
[:=] A \vdash [x'' = -g] \left((x = 0 \rightarrow [v := -cv] B(x, v)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
\hline
[?] A \vdash [x'' = -g] \left([?x = 0] [v := -cv] B(x, v) \wedge [?x \geq 0] B(x, v) \right) \\
\hline
[.] A \vdash [x'' = -g] \left([?x = 0; v := -cv] B(x, v) \wedge [?x \geq 0] B(x, v) \right) \\
\hline
[U] A \vdash [x'' = -g] [?x = 0; v := -cv \cup ?x \geq 0] B(x, v) \\
\hline
[.] A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)] B(x, v) \\
\hline
\rightarrow^R \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)] B(x, v)
\end{array}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$$[x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v) \leftrightarrow$$

$$[x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v) \text{ by } [;]$$

$$\frac{[U] \frac{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}{[;] A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}}{\rightarrow^R \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$$[?x = 0; v := -cv \cup ?x \geq 0]B(x, v) \leftrightarrow$$

$$([?x = 0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v)) \text{ by } [\cup]$$

$$\frac{[i] \frac{A \vdash [x'' = -g]([?x = 0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}{[U] A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}}{[i] A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}}{\rightarrow^R \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$$[?x = 0; v := -cv]B(x, v) \leftrightarrow$$

$$[?x = 0][v := -cv]B(x, v) \text{ by } [;]$$

$$\frac{[?]}{A \vdash [x'' = -g]([?x = 0][v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}$$

$$\frac{[;]}{A \vdash [x'' = -g]([?x = 0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}$$

$$\frac{[U]}{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}$$

$$\frac{[;]}{A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$\rightarrow^R \frac{}{\vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$[?x = 0][v := -cv]B(x, v) \leftrightarrow$
 $x = 0 \rightarrow [v := -cv]B(x, v)$ by [?]

$$\frac{[:=]}{A \vdash [x'' = -g]((x = 0 \rightarrow [v := -cv]B(x, v)) \wedge (x \geq 0 \rightarrow B(x, v)))}$$

$$\frac{[?]}{A \vdash [x'' = -g]([?x = 0][v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}$$

$$\frac{[;]}{A \vdash [x'' = -g]([?x = 0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}$$

$$\frac{[U]}{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}$$

$$\frac{[;]}{A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$\frac{\rightarrow R}{\vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$$[v := -cv]B(x, v) \leftrightarrow \\ x = 0 \rightarrow B(x, -cv) \text{ by } [:=]$$

$$\begin{array}{c} \frac{[?]}{A \vdash [x'' = -g]((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))} \\ \frac{[:=]}{A \vdash [x'' = -g]((x = 0 \rightarrow [v := -cv]B(x, v)) \wedge (x \geq 0 \rightarrow B(x, v)))} \\ \frac{[?]}{A \vdash [x'' = -g]([?x = 0][v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))} \\ \frac{[?]}{A \vdash [x'' = -g]([?x = 0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))} \\ \frac{[U]}{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)} \\ \frac{[?]}{A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)} \\ \rightarrow^R \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v) \end{array}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$$[\cdot] \quad [x' = f(x)]p(x) \leftrightarrow \forall t \geq 0 [x := y(t)]p(x)$$

$$[\cdot] \quad \frac{}{A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2; v := -gt] ((x=0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))}$$

$$[\cdot] \quad \frac{}{A \vdash [x'' = -g] ((x=0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))}$$

$$[\cdot :=] \quad \frac{}{A \vdash [x'' = -g] ((x=0 \rightarrow [v := -cv]B(x, v)) \wedge (x \geq 0 \rightarrow B(x, v)))}$$

$$[\cdot ?] \quad \frac{}{A \vdash [x'' = -g] ([?x=0][v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}$$

$$[\cdot] \quad \frac{}{A \vdash [x'' = -g] ([?x=0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}$$

$$[\cdot \cup] \quad \frac{}{A \vdash [x'' = -g][?x=0; v := -cv \cup ?x \geq 0]B(x, v)}$$

$$[\cdot] \quad \frac{}{A \vdash [x'' = -g; (?x=0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$\rightarrow^R \quad \frac{}{\vdash A \rightarrow [x'' = -g; (?x=0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$$\frac{[]}{A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2][v := -gt] ((x=0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))}$$

$$\frac{[]}{A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2; v := -gt] ((x=0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))}$$

$$\frac{[]}{A \vdash [x'' = -g] ((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))}$$

$$\frac{[]}{A \vdash [x'' = -g] ((x = 0 \rightarrow [v := -cv]B(x, v)) \wedge (x \geq 0 \rightarrow B(x, v)))}$$

$$\frac{[]}{A \vdash [x'' = -g] ([?x = 0][v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}$$

$$\frac{[]}{A \vdash [x'' = -g] ([?x = 0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}$$

$$\frac{[]}{A \vdash [x'' = -g][?x = 0; v := -cv \cup ?x \geq 0]B(x, v)}$$

$$\frac{[]}{A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)}$$

$$\rightarrow^R \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)]B(x, v)$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

$$\frac{[]}{[:=]} \frac{A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2] ((x=0 \rightarrow B(x, -c(-gt))) \wedge (x \geq 0 \rightarrow B(x, -gt)))}{[:=]} \frac{A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2] [v := -gt] ((x=0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))}{[:=]} \frac{A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2; v := -gt] ((x=0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))}{[:=]} \frac{A \vdash [x'' = -g] ((x=0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)))}{[:=]} \frac{A \vdash [x'' = -g] ((x=0 \rightarrow [v := -cv]B(x, v)) \wedge (x \geq 0 \rightarrow B(x, v)))}{[:=]} \frac{A \vdash [x'' = -g] ([?x=0][v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}{[?]} \frac{A \vdash [x'' = -g] ([?x=0; v := -cv]B(x, v) \wedge [?x \geq 0]B(x, v))}{[:=]} \frac{A \vdash [x'' = -g][?x=0; v := -cv \cup ?x \geq 0]B(x, v)}{[U]} \frac{A \vdash [x'' = -g; (?x=0; v := -cv \cup ?x \geq 0)]B(x, v)}{[:=]} \frac{\rightarrow^R}{\vdash} \frac{A \rightarrow [x'' = -g; (?x=0; v := -cv \cup ?x \geq 0)]B(x, v)}{A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$



A Sequent Proof of a Single-hop Bouncing Ball



$$\begin{array}{c}
 A \vdash \forall t \geq 0 \left(\left(H - \frac{g}{2}t^2 = 0 \rightarrow B\left(H - \frac{g}{2}t^2, -c(-gt)\right) \right) \wedge \left(H - \frac{g}{2}t^2 \geq 0 \rightarrow B\left(H - \frac{g}{2}t^2, -g\right) \right) \right) \\
 \hline
 [:=] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2] \left((x = 0 \rightarrow B(x, -c(-gt))) \wedge (x \geq 0 \rightarrow B(x, -gt)) \right) \\
 \hline
 [:=] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2] [v := -gt] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
 \hline
 [:] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2; v := -gt] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
 \hline
 ['] A \vdash [x'' = -g] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
 \hline
 [:=] A \vdash [x'' = -g] \left((x = 0 \rightarrow [v := -cv] B(x, v)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
 \hline
 [?] A \vdash [x'' = -g] \left([?x = 0] [v := -cv] B(x, v) \wedge [?x \geq 0] B(x, v) \right) \\
 \hline
 [:] A \vdash [x'' = -g] \left([?x = 0; v := -cv] B(x, v) \wedge [?x \geq 0] B(x, v) \right) \\
 \hline
 [U] A \vdash [x'' = -g] [?x = 0; v := -cv \cup ?x \geq 0] B(x, v) \\
 \hline
 [:] A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)] B(x, v) \\
 \hline
 \rightarrow^R \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)] B(x, v)
 \end{array}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$



A Sequent Proof of a Single-hop Bouncing Ball



$$\begin{array}{c}
 A \vdash \forall t \geq 0 \left(\left(H - \frac{g}{2}t^2 = 0 \rightarrow B\left(H - \frac{g}{2}t^2, -c(-gt)\right) \right) \wedge \left(H - \frac{g}{2}t^2 \geq 0 \rightarrow B\left(H - \frac{g}{2}t^2, -g\right) \right) \right) \\
 \hline
 [:=] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2] \left((x = 0 \rightarrow B(x, -c(-gt))) \wedge (x \geq 0 \rightarrow B(x, -gt)) \right) \\
 \hline
 [:=] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2] [v := -gt] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
 \hline
 [:] A \vdash \forall t \geq 0 [x := H - \frac{g}{2}t^2; v := -gt] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
 \hline
 ['] A \vdash [x'' = -g] \left((x = 0 \rightarrow B(x, -cv)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
 \hline
 [:=] A \vdash [x'' = -g] \left((x = 0 \rightarrow [v := -cv] B(x, v)) \wedge (x \geq 0 \rightarrow B(x, v)) \right) \\
 \hline
 [?] A \vdash [x'' = -g] \left([?x = 0] [v := -cv] B(x, v) \wedge [?x \geq 0] B(x, v) \right) \\
 \hline
 [:] A \vdash [x'' = -g] \left([?x = 0; v := -cv] B(x, v) \wedge [?x \geq 0] B(x, v) \right) \\
 \hline
 [U] A \vdash [x'' = -g] [?x = 0; v := -cv \cup ?x \geq 0] B(x, v) \\
 \hline
 [:] A \vdash [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)] B(x, v) \\
 \hline
 \rightarrow^R \vdash A \rightarrow [x'' = -g; (?x = 0; v := -cv \cup ?x \geq 0)] B(x, v)
 \end{array}$$

$$A \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x = H \wedge v = 0 \wedge g > 0 \wedge 1 \geq c \geq 0$$

$$B(x, v) \stackrel{\text{def}}{\equiv} 0 \leq x \wedge x \leq H$$

$$\{x'' = -g\} \stackrel{\text{def}}{\equiv} \{x' = v, v' = -g\}$$

1 Learning Objectives

2 Sequent Calculus

- Propositional Proof Rules
- Soundness of Proof Rules
- Proofs with Dynamics
- Contextual Equivalence
- Quantifier Proof Rules
- A Sequent Proof for Single-hop Bouncing Balls

3 Real Arithmetic

- Real Quantifier Elimination
- Instantiating Real-Arithmetic Quantifiers
- Weakening by Removing Assumptions
- Abbreviating Terms to Reduce Complexity
- Substituting Equations into Formulas
- Creatively Cutting to Transform Questions

4 Summary

Lemma (\mathbb{R} real arithmetic)

$\text{FOL}_{\mathbb{R}}$ decidable, so side condition implementable:

$$\mathbb{R} \frac{}{\Gamma \vdash \Delta} \quad \left(\text{if } \bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q \text{ is valid in } \text{FOL}_{\mathbb{R}} \right)$$

$$\overline{\mathbb{R} a > 0, b > 0 \vdash y \geq 0 \rightarrow ax^2 + by \geq 0}$$

$$\overline{\mathbb{R} x^2 > 0 \vdash x > 0}$$

Lemma (\mathbb{R} real arithmetic)

$\text{FOL}_{\mathbb{R}}$ decidable, so side condition implementable:

$$\mathbb{R} \frac{}{\Gamma \vdash \Delta} \quad \left(\text{if } \bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q \text{ is valid in } \text{FOL}_{\mathbb{R}} \right)$$

*

$$\mathbb{R} \frac{}{a > 0, b > 0 \vdash y \geq 0 \rightarrow ax^2 + by \geq 0}$$

$$\mathbb{R} \frac{}{x^2 > 0 \vdash x > 0}$$

Lemma (\mathbb{R} real arithmetic)

$\text{FOL}_{\mathbb{R}}$ decidable, so side condition implementable:

$$\mathbb{R} \frac{}{\Gamma \vdash \Delta} \quad \left(\text{if } \bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q \text{ is valid in } \text{FOL}_{\mathbb{R}} \right)$$

$$\mathbb{R} \frac{*}{a > 0, b > 0 \vdash y \geq 0 \rightarrow ax^2 + by \geq 0}$$

$$\mathbb{R} \frac{\text{false}}{x^2 > 0 \vdash x > 0}$$

Lemma (\mathbb{R} real arithmetic)

$\text{FOL}_{\mathbb{R}}$ *decidable, so side condition implementable:*

$$\mathbb{R} \frac{}{\Gamma \vdash \Delta} \quad \left(\text{if } \bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q \text{ is valid in } \text{FOL}_{\mathbb{R}} \right)$$

$$\mathbb{R} \frac{*}{a > 0, b > 0 \vdash y \geq 0 \rightarrow ax^2 + by \geq 0}$$

$$\mathbb{R} \frac{\text{false}}{x^2 > 0 \vdash x > 0}$$

Theorem (Tarski's quantifier elimination)

$\text{FOL}_{\mathbb{R}}$ *admits quantifier elimination: there is an algorithm that computes a quantifier-free formula $\text{QE}(P)$, for each first-order real arithmetic formula P , that is equivalent, i.e., $P \leftrightarrow \text{QE}(P)$ is valid.*

Lemma (\mathbb{R} real arithmetic)

$\text{FOL}_{\mathbb{R}}$ *decidable, so side condition implementable:*

$$\mathbb{R} \frac{}{\Gamma \vdash \Delta} \quad \left(\text{if } \bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q \text{ is valid in } \text{FOL}_{\mathbb{R}} \right)$$

$$\mathbb{R} \frac{*}{a > 0, b > 0 \vdash y \geq 0 \rightarrow ax^2 + by \geq 0}$$

$$\mathbb{R} \frac{\text{false}}{x^2 > 0 \vdash x > 0}$$

Theorem (Tarski's quantifier elimination)

$\text{FOL}_{\mathbb{R}}$ *admits quantifier elimination: there is an algorithm that computes a quantifier-free formula $\text{QE}(P)$, for each first-order real arithmetic formula P , that is equivalent, i.e., $P \leftrightarrow \text{QE}(P)$ is valid.*

What if there are no quantifiers?

Lemma (\mathbb{R} real arithmetic)

$\text{FOL}_{\mathbb{R}}$ decidable, so side condition implementable:

$$\mathbb{R} \frac{}{\Gamma \vdash \Delta} \quad \left(\text{if } \bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q \text{ is valid in } \text{FOL}_{\mathbb{R}} \right)$$

$$\mathbb{R} \frac{*}{a > 0, b > 0 \vdash y \geq 0 \rightarrow ax^2 + by \geq 0}$$

$$\mathbb{R} \frac{\text{false}}{x^2 > 0 \vdash x > 0}$$

Theorem (Tarski's quantifier elimination)

$\text{FOL}_{\mathbb{R}}$ admits quantifier elimination: there is an algorithm that computes a quantifier-free formula $\text{QE}(P)$, for each first-order real arithmetic formula P , that is equivalent, i.e., $P \leftrightarrow \text{QE}(P)$ is valid.

What if there are no quantifiers? Universal closure with \forall $\frac{\Gamma \vdash \forall x P, \Delta}{\Gamma \vdash P, \Delta}$



$$\forall R \frac{}{\vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)}$$



$$\forall^{\mathbb{R}} \overline{\vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)}$$

Not a $\text{FOL}_{\mathbb{R}}$ formula so Tarski's quantifier elimination not applicable.



$$\frac{[U] \quad \vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0}{\forall R \quad \vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)}$$

$$\begin{array}{c}
 \frac{}{[:=] \vdash d \geq -x \rightarrow [x := 0] x \geq 0 \wedge [x := x + d] x \geq 0} \\
 \frac{}{[U] \vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0} \\
 \frac{}{\forall R \vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)}
 \end{array}$$

$$\begin{array}{c}
 \frac{}{[:=] \vdash d \geq -x \rightarrow 0 \geq 0 \wedge [x := x + d] x \geq 0} \\
 \frac{}{[:=] \vdash d \geq -x \rightarrow [x := 0] x \geq 0 \wedge [x := x + d] x \geq 0} \\
 \frac{}{[U] \vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0} \\
 \frac{}{\forall R \vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)}
 \end{array}$$



$$\begin{array}{c} \text{i}\forall \frac{}{\vdash d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0} \\ \text{[:=]} \frac{}{\vdash d \geq -x \rightarrow 0 \geq 0 \wedge [x := x + d] x \geq 0} \\ \text{[:=]} \frac{}{\vdash d \geq -x \rightarrow [x := 0] x \geq 0 \wedge [x := x + d] x \geq 0} \\ \text{[U]} \frac{}{\vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0} \\ \forall R \frac{}{\vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)} \end{array}$$



$$\begin{array}{c} \text{i}\forall \frac{}{\vdash \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0)} \\ \text{i}\forall \frac{}{\vdash d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0} \\ [\text{:=}] \frac{}{\vdash d \geq -x \rightarrow 0 \geq 0 \wedge [x := x + d] x \geq 0} \\ [\text{:=}] \frac{}{\vdash d \geq -x \rightarrow [x := 0] x \geq 0 \wedge [x := x + d] x \geq 0} \\ [\cup] \frac{}{\vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0} \\ \forall R \frac{}{\vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)} \end{array}$$

$$\begin{array}{l}
\mathbb{R} \frac{}{\vdash \forall x \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0)} \\
\text{i}\forall \frac{}{\vdash \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0)} \\
\text{i}\forall \frac{}{\vdash d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0} \\
[:=] \frac{}{\vdash d \geq -x \rightarrow 0 \geq 0 \wedge [x := x + d] x \geq 0} \\
[:=] \frac{}{\vdash d \geq -x \rightarrow [x := 0] x \geq 0 \wedge [x := x + d] x \geq 0} \\
[\cup] \frac{}{\vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0} \\
\forall\mathbb{R} \frac{}{\vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)}
\end{array}$$

$$\begin{array}{c}
 * \\
 \hline
 \mathbb{R} \quad \vdash \forall x \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0) \\
 \hline
 \text{i}\forall \quad \vdash \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0) \\
 \hline
 \text{i}\forall \quad \vdash d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0 \\
 \hline
 [:=] \quad \vdash d \geq -x \rightarrow 0 \geq 0 \wedge [x := x + d] x \geq 0 \\
 \hline
 [:=] \quad \vdash d \geq -x \rightarrow [x := 0] x \geq 0 \wedge [x := x + d] x \geq 0 \\
 \hline
 [\cup] \quad \vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0 \\
 \hline
 \forall\mathbb{R} \quad \vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)
 \end{array}$$

$$\begin{array}{c}
 * \\
 \hline
 \mathbb{R} \vdash \forall x \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0) \\
 \hline
 \text{i}\forall \vdash \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0) \\
 \hline
 \text{ii}\forall \vdash d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0 \\
 \hline
 [:=] \vdash d \geq -x \rightarrow 0 \geq 0 \wedge [x := x + d] x \geq 0 \\
 \hline
 [:=] \vdash d \geq -x \rightarrow [x := 0] x \geq 0 \wedge [x := x + d] x \geq 0 \\
 \hline
 [\cup] \vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0 \\
 \hline
 \forall\mathbb{R} \vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)
 \end{array}$$

We could also leave $\forall d$ alone and use axioms in the middle of the formula.

$$\begin{array}{c}
 * \\
 \mathbb{R} \frac{}{\vdash \forall x \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0)} \\
 i\forall \frac{}{\vdash \forall d (d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0)} \\
 i\forall \frac{}{\vdash d \geq -x \rightarrow 0 \geq 0 \wedge x + d \geq 0} \\
 [:=] \frac{}{\vdash d \geq -x \rightarrow 0 \geq 0 \wedge [x := x + d] x \geq 0} \\
 [:=] \frac{}{\vdash d \geq -x \rightarrow [x := 0] x \geq 0 \wedge [x := x + d] x \geq 0} \\
 [U] \frac{}{\vdash d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0} \\
 \forall\mathbb{R} \frac{}{\vdash \forall d (d \geq -x \rightarrow [x := 0 \cup x := x + d] x \geq 0)}
 \end{array}$$

Already use rule \mathbb{R} for valid $\text{FOL}_{\mathbb{R}}$ formulas with free variables before $i\forall$



Instantiating Real-Arithmetic Quantifiers

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\Gamma \vdash [x' = f(x) \& q(x)]P$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\frac{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P)}{\Gamma \vdash [x' = f(x) \& q(x)]P}$$



Instantiating Real-Arithmetic Quantifiers

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\frac{\frac{\frac{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s)))) \rightarrow [x := y(t)] P}{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s)))) \rightarrow [x := y(t)] P}}{[\Gamma]} \Gamma \vdash [x' = f(x) \& q(x)] P$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\frac{\frac{\frac{\Gamma, t \geq 0 \vdash (\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)] P}{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s))) \rightarrow [x := y(t)] P)}}{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)] P)}}{[\cdot]} \Gamma \vdash [x' = f(x) \& q(x)] P$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\begin{array}{l} \frac{}{\rightarrow R} \frac{\Gamma, t \geq 0, \forall 0 \leq s \leq t q(y(s)) \vdash [x := y(t)]P}{\Gamma, t \geq 0 \vdash (\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P} \\ \frac{}{\rightarrow R} \frac{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s))) \rightarrow [x := y(t)]P)}{\forall R} \frac{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P)}{[\cdot]} \frac{}{\Gamma \vdash [x' = f(x) \& q(x)]P} \end{array}$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\frac{}{\forall L \frac{\Gamma, t \geq 0, 0 \leq t \leq t \rightarrow q(y(t)) \vdash [x := y(t)]P}{\Gamma, t \geq 0, \forall 0 \leq s \leq t q(y(s)) \vdash [x := y(t)]P}} \rightarrow R \frac{}{\rightarrow R \frac{\Gamma, t \geq 0 \vdash (\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P}{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s)))) \rightarrow [x := y(t)]P}} \forall R \frac{}{[\cdot] \frac{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s)))) \rightarrow [x := y(t)]P}{\Gamma \vdash [x' = f(x) \& q(x)]P}}$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\begin{array}{l} \rightarrow L \frac{\overline{t \geq 0 \vdash 0 \leq t \leq t, [x := y(t)]P} \quad \overline{\Gamma, t \geq 0, q(y(t)) \vdash [x := y(t)]P}}{\Gamma, t \geq 0, 0 \leq t \leq t \rightarrow q(y(t)) \vdash [x := y(t)]P} \\ \forall L \frac{\Gamma, t \geq 0, 0 \leq t \leq t \rightarrow q(y(t)) \vdash [x := y(t)]P}{\Gamma, t \geq 0, \forall 0 \leq s \leq t q(y(s)) \vdash [x := y(t)]P} \\ \rightarrow R \frac{\Gamma, t \geq 0, \forall 0 \leq s \leq t q(y(s)) \vdash [x := y(t)]P}{\Gamma, t \geq 0 \vdash (\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P} \\ \rightarrow R \frac{\Gamma, t \geq 0 \vdash (\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P}{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s)))) \rightarrow [x := y(t)]P} \\ \forall R \frac{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s)))) \rightarrow [x := y(t)]P}{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s)))) \rightarrow [x := y(t)]P} \\ [\cdot] \frac{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s)))) \rightarrow [x := y(t)]P}{\Gamma \vdash [x' = f(x) \& q(x)]P} \end{array}$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\begin{array}{c}
 * \\
 \mathbb{R} \frac{}{t \geq 0 \vdash 0 \leq t \leq t, [x := y(t)]P} \quad \frac{}{\Gamma, t \geq 0, q(y(t)) \vdash [x := y(t)]P} \\
 \rightarrow L \frac{}{\Gamma, t \geq 0, 0 \leq t \leq t \rightarrow q(y(t)) \vdash [x := y(t)]P} \\
 \forall L \frac{}{\Gamma, t \geq 0, \forall 0 \leq s \leq t q(y(s)) \vdash [x := y(t)]P} \\
 \rightarrow R \frac{}{\Gamma, t \geq 0 \vdash (\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P} \\
 \rightarrow R \frac{}{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s))) \rightarrow [x := y(t)]P)} \\
 \forall R \frac{}{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P)} \\
 [\cdot] \frac{}{\Gamma \vdash [x' = f(x) \& q(x)]P}
 \end{array}$$

$$\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (\dots) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\dots)$$

$$\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\dots) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (\dots)$$

$$\begin{array}{c} * \qquad \qquad \qquad \dots \\ \mathbb{R} \frac{}{t \geq 0 \vdash 0 \leq t \leq t, [x := y(t)]P} \quad \frac{}{\Gamma, t \geq 0, q(y(t)) \vdash [x := y(t)]P} \\ \rightarrow L \frac{}{\Gamma, t \geq 0, 0 \leq t \leq t \rightarrow q(y(t)) \vdash [x := y(t)]P} \\ \forall L \frac{}{\Gamma, t \geq 0, \forall 0 \leq s \leq t q(y(s)) \vdash [x := y(t)]P} \\ \rightarrow R \frac{}{\Gamma, t \geq 0 \vdash (\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P} \\ \rightarrow R \frac{}{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s))) \rightarrow [x := y(t)]P)} \\ \forall R \frac{}{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P)} \\ [\cdot] \frac{}{\Gamma \vdash [x' = f(x) \& q(x)]P} \end{array}$$

Derived Rule

$$\frac{\Gamma, t \geq 0, q(y(t)) \vdash [x := y(t)]P}{\Gamma \vdash [x' = f(x) \& q(x)]P} \quad (y'(t) = f(y))$$

$$\frac{\begin{array}{c} \mathbb{R} \\ \frac{t \geq 0 \vdash 0 \leq t \leq t, [x := y(t)]P}{\rightarrow L} \quad \frac{\Gamma, t \geq 0, q(y(t)) \vdash [x := y(t)]P}{\dots} \\ \frac{\Gamma, t \geq 0, 0 \leq t \leq t \rightarrow q(y(t)) \vdash [x := y(t)]P}{\forall L} \\ \frac{\Gamma, t \geq 0, \forall 0 \leq s \leq t q(y(s)) \vdash [x := y(t)]P}{\rightarrow R} \\ \frac{\Gamma, t \geq 0 \vdash (\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P}{\rightarrow R} \\ \frac{\Gamma \vdash t \geq 0 \rightarrow ((\forall 0 \leq s \leq t Q(y(s))) \rightarrow [x := y(t)]P)}{\forall R} \\ \frac{\Gamma \vdash \forall t \geq 0 ((\forall 0 \leq s \leq t q(y(s))) \rightarrow [x := y(t)]P)}{[']} \\ \Gamma \vdash [x' = f(x) \& q(x)]P \end{array}}{\Gamma \vdash [x' = f(x) \& q(x)]P}$$

Derived rule: rule that can be proved using other proof rules.

$$\text{WR} \frac{\Gamma \vdash \Delta}{\Gamma \vdash P, \Delta}$$

$$\text{WL} \frac{\Gamma \vdash \Delta}{\Gamma, P \vdash \Delta}$$

$$\text{WL} \frac{r \geq 0 \vdash 0 \leq r \leq r}{A, r \geq 0 \vdash 0 \leq r \leq r}$$

Throw big arithmetic distraction A away by weakening since the proof is independent of formula A .

Occam's assumption razor

Think how hard it would be to prove a theorem with all the facts in all books of mathematics as assumptions.

Compared to a proof from just the two facts that matter.



$$a \geq 0, t \geq 0, 0 \leq \underbrace{\frac{a}{2}t^2 + vt + x}_z, \underbrace{\frac{a}{2}t^2 + vt + x}_z \leq d, d \leq 8 \vdash \underbrace{\frac{a}{2}t^2 + vt + x}_z \leq 8$$

Abbreviate fancy term $\frac{a}{2}t^2 + vt + x$ by new variable z makes it easy:

$$a \geq 0, t \geq 0, 0 \leq z, z \leq d, d \leq 8 \vdash z \leq 8$$



$$a \geq 0, t \geq 0, 0 \leq \underbrace{\frac{a}{2}t^2 + vt + x}_z, \underbrace{\frac{a}{2}t^2 + vt + x}_z \leq d, d \leq 8 \vdash \underbrace{\frac{a}{2}t^2 + vt + x}_z \leq 8$$

Abbreviate fancy term $\frac{a}{2}t^2 + vt + x$ by new variable z makes it easy:

$$a \geq 0, t \geq 0, 0 \leq z, z \leq d, d \leq 8 \vdash z \leq 8$$

Proof rules introducing such new variables will be studied in [Chapter 12](#)



$$a \geq 0, t \geq 0, 0 \leq \underbrace{\frac{a}{2}t^2 + vt + x}_z, \underbrace{\frac{a}{2}t^2 + vt + x}_z \leq d, d \leq 8 \vdash \underbrace{\frac{a}{2}t^2 + vt + x}_z \leq 8$$

Abbreviate fancy term $\frac{a}{2}t^2 + vt + x$ by new variable z makes it easy:

$$a \geq 0, t \geq 0, 0 \leq z, z \leq d, d \leq 8 \vdash z \leq 8$$

Proof rules introducing such new variables will be studied in [Chapter 12](#)

Inverse of a derived rule that turns assignments into equations:

$$[:=] = \frac{\Gamma, y = e \vdash p(y), \Delta}{\Gamma \vdash [x := e]p(x), \Delta}$$



$$a \geq 0, t \geq 0, 0 \leq \underbrace{\frac{a}{2}t^2 + vt + x}_z, \underbrace{\frac{a}{2}t^2 + vt + x}_z \leq d, d \leq 8 \vdash \underbrace{\frac{a}{2}t^2 + vt + x}_z \leq 8$$

Abbreviate fancy term $\frac{a}{2}t^2 + vt + x$ by new variable z makes it easy:

$$a \geq 0, t \geq 0, 0 \leq z, z \leq d, d \leq 8 \vdash z \leq 8$$

Proof rules introducing such new variables will be studied in [Chapter 12](#)

Inverse of a derived rule that turns assignments into equations:

$$[:=] = \frac{\Gamma, y = e \vdash p(y), \Delta}{\Gamma \vdash [x := e]p(x), \Delta} \quad (y \text{ new})$$



$$=R \frac{\Gamma, x = e \vdash p(e), \Delta}{\Gamma, x = e \vdash p(x), \Delta}$$

$$=L \frac{\Gamma, x = e, p(e) \vdash \Delta}{\Gamma, x = e, p(x) \vdash \Delta}$$

$$\begin{array}{c} \text{cut} \\ \hline (x-y)^2 \leq 0, p(y) \vdash p(x) \\ \hline \wedge L \\ (x-y)^2 \leq 0 \wedge p(y) \vdash p(x) \\ \hline \rightarrow R \\ \vdash (x-y)^2 \leq 0 \wedge p(y) \rightarrow p(x) \end{array}$$



$$=R \frac{\Gamma, x = e \vdash p(e), \Delta}{\Gamma, x = e \vdash p(x), \Delta}$$

$$=L \frac{\Gamma, x = e, p(e) \vdash \Delta}{\Gamma, x = e, p(x) \vdash \Delta}$$

$$\begin{array}{c} \text{cut} \\ \frac{\frac{\text{WL} \overline{(x-y)^2 \leq 0, p(y) \vdash x = y, p(x)} \quad \text{WL} \overline{(x-y)^2 \leq 0, p(y), x = y \vdash p(x)}}{(x-y)^2 \leq 0, p(y) \vdash p(x)}}{\text{WL} \overline{(x-y)^2 \leq 0 \wedge p(y) \vdash p(x)}} \\ \text{AND} \\ \frac{\text{WL} \overline{(x-y)^2 \leq 0 \wedge p(y) \vdash p(x)}}{\text{WL} \overline{\vdash (x-y)^2 \leq 0 \wedge p(y) \rightarrow p(x)}} \\ \text{IMPL} \end{array}$$



Creatively Cutting to Transform Questions

$$=R \frac{\Gamma, x = e \vdash p(e), \Delta}{\Gamma, x = e \vdash p(x), \Delta}$$

$$=L \frac{\Gamma, x = e, p(e) \vdash \Delta}{\Gamma, x = e, p(x) \vdash \Delta}$$

$$\begin{array}{c}
 * \\
 \mathbb{R} \frac{}{(x-y)^2 \leq 0 \vdash x = y} \\
 WR \frac{}{(x-y)^2 \leq 0 \vdash x = y, p(x)} \\
 WL \frac{}{(x-y)^2 \leq 0, p(y) \vdash x = y, p(x)} \quad WL \frac{}{(x-y)^2 \leq 0, p(y), x = y \vdash p(x)} \\
 cut \frac{}{(x-y)^2 \leq 0, p(y) \vdash p(x)} \\
 \wedge L \frac{}{(x-y)^2 \leq 0 \wedge p(y) \vdash p(x)} \\
 \rightarrow R \frac{}{\vdash (x-y)^2 \leq 0 \wedge p(y) \rightarrow p(x)}
 \end{array}$$

$$\begin{aligned}
 &=R \frac{\Gamma, x = e \vdash p(e), \Delta}{\Gamma, x = e \vdash p(x), \Delta} \\
 &=L \frac{\Gamma, x = e, p(e) \vdash \Delta}{\Gamma, x = e, p(x) \vdash \Delta}
 \end{aligned}$$

$$\begin{array}{c}
 * \\
 \mathbb{R} \frac{}{(x-y)^2 \leq 0 \vdash x = y} \\
 WR \frac{}{(x-y)^2 \leq 0 \vdash x = y, p(x)} \\
 WL \frac{}{(x-y)^2 \leq 0, p(y) \vdash x = y, p(x)} \\
 \text{cut} \frac{}{(x-y)^2 \leq 0, p(y) \vdash p(x)} \\
 \wedge L \frac{}{(x-y)^2 \leq 0 \wedge p(y) \vdash p(x)} \\
 \rightarrow R \frac{}{\vdash (x-y)^2 \leq 0 \wedge p(y) \rightarrow p(x)}
 \end{array}
 \quad
 \begin{array}{c}
 =R \frac{}{p(y), x = y \vdash p(x)} \\
 WL \frac{}{(x-y)^2 \leq 0, p(y), x = y \vdash p(x)}
 \end{array}$$



$$\begin{aligned}
 &=R \frac{\Gamma, x = e \vdash p(e), \Delta}{\Gamma, x = e \vdash p(x), \Delta} \\
 &=L \frac{\Gamma, x = e, p(e) \vdash \Delta}{\Gamma, x = e, p(x) \vdash \Delta}
 \end{aligned}$$

$$\begin{array}{c}
 * \\
 \mathbb{R} \frac{}{(x-y)^2 \leq 0 \vdash x = y} \\
 WR \frac{}{(x-y)^2 \leq 0 \vdash x = y, p(x)} \\
 WL \frac{}{(x-y)^2 \leq 0, p(y) \vdash x = y, p(x)} \\
 \text{cut} \frac{}{(x-y)^2 \leq 0, p(y) \vdash p(x)} \\
 \wedge L \frac{}{(x-y)^2 \leq 0 \wedge p(y) \vdash p(x)} \\
 \rightarrow R \frac{}{\vdash (x-y)^2 \leq 0 \wedge p(y) \rightarrow p(x)}
 \end{array}
 \quad
 \begin{array}{c}
 \text{id} \frac{}{p(y), x = y \vdash p(y)} \\
 =R \frac{}{p(y), x = y \vdash p(x)} \\
 WL \frac{}{(x-y)^2 \leq 0, p(y), x = y \vdash p(x)}
 \end{array}$$



$$\begin{aligned}
 &=R \frac{\Gamma, x = e \vdash p(e), \Delta}{\Gamma, x = e \vdash p(x), \Delta} \\
 &=L \frac{\Gamma, x = e, p(e) \vdash \Delta}{\Gamma, x = e, p(x) \vdash \Delta}
 \end{aligned}$$

| | |
|---|--|
| * | * |
| $\mathbb{R} \frac{}{(x-y)^2 \leq 0 \vdash x = y}$ | $\text{id} \frac{}{p(y), x = y \vdash p(y)}$ |
| $\text{WR} \frac{}{(x-y)^2 \leq 0 \vdash x = y, p(x)}$ | $=R \frac{}{p(y), x = y \vdash p(x)}$ |
| $\text{WL} \frac{}{(x-y)^2 \leq 0, p(y) \vdash x = y, p(x)}$ | $\text{WL} \frac{}{(x-y)^2 \leq 0, p(y), x = y \vdash p(x)}$ |
| $\text{cut} \frac{}{(x-y)^2 \leq 0, p(y) \vdash p(x)}$ | |
| $\wedge L \frac{}{(x-y)^2 \leq 0 \wedge p(y) \vdash p(x)}$ | |
| $\rightarrow R \frac{}{\vdash (x-y)^2 \leq 0 \wedge p(y) \rightarrow p(x)}$ | |



1 Learning Objectives

2 Sequent Calculus

- Propositional Proof Rules
- Soundness of Proof Rules
- Proofs with Dynamics
- Contextual Equivalence
- Quantifier Proof Rules
- A Sequent Proof for Single-hop Bouncing Balls

3 Real Arithmetic

- Real Quantifier Elimination
- Instantiating Real-Arithmetic Quantifiers
- Weakening by Removing Assumptions
- Abbreviating Terms to Reduce Complexity
- Substituting Equations into Formulas
- Creatively Cutting to Transform Questions

4 Summary

Summary: Proof Rules of Sequent Calculus

$$\begin{array}{c}
 \neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta} \quad \wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta} \quad \vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta} \\
 \neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta} \quad \wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta} \quad \vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta} \\
 \rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta} \quad \text{id} \frac{}{\Gamma, P \vdash P, \Delta} \quad \top R \frac{}{\Gamma \vdash \text{true}, \Delta} \\
 \rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta} \quad \text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta} \quad \perp L \frac{}{\Gamma, \text{false} \vdash \Delta} \\
 \forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (y \notin \Gamma, \Delta, \forall x p(x)) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\text{arbitrary term } e) \\
 \forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\text{arbitrary term } e) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (y \notin \Gamma, \Delta, \exists x p(x))
 \end{array}$$

Summary: Proof Rules of Sequent Calculus

$$\begin{array}{c}
 \neg R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \neg P, \Delta} \quad \wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta} \quad \vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta} \\
 \neg L \frac{\Gamma \vdash P, \Delta}{\Gamma, \neg P \vdash \Delta} \quad \wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta} \quad \vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta} \\
 \rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta} \quad \text{id} \frac{}{\Gamma, P \vdash P, \Delta} \quad \text{TR} \frac{}{\Gamma \vdash \text{true}, \Delta} \\
 \rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta} \quad \text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta} \quad \perp L \frac{}{\Gamma, \text{false} \vdash \Delta} \\
 \forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} (y \notin \Gamma, \Delta, \forall x p(x)) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} (\text{arbitrary term } e) \\
 \forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} (\text{arbitrary term } e) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} (y \notin \Gamma, \Delta, \exists x p(x)) \\
 \mathbb{R} \frac{}{\Gamma \vdash \Delta} \quad (\text{if } \bigwedge_{P \in \Gamma} P \rightarrow \bigvee_{Q \in \Delta} Q \text{ is valid in } \text{FOL}_{\mathbb{R}})
 \end{array}$$



André Platzer.

Logical Foundations of Cyber-Physical Systems.

Springer, Cham, 2018.

doi:10.1007/978-3-319-63588-0.



André Platzer.

Logical Analysis of Hybrid Systems: Proving Theorems for Complex Dynamics.

Springer, Heidelberg, 2010.

doi:10.1007/978-3-642-14509-4.



André Platzer.

Differential dynamic logic for hybrid systems.

J. Autom. Reas., 41(2):143–189, 2008.

doi:10.1007/s10817-008-9103-8.